

Controlled Substitution of Soy Protein for Meat Protein: Effects on Calcium Retention, Bone, and Cardiovascular Health Indices in Postmenopausal Women

Zamzam K. (Fariba) Roughead, Janet R. Hunt, LuAnn K. Johnson, Thomas M. Badger, and Glenn I. Lykken

United States Department of Agriculture (Z.K.R., J.R.H., L.K.J.), Agricultural Research Service, Grand Forks Human Nutrition Research Center, Grand Forks, North Dakota 58202-9034; Arkansas Children's Nutrition Center (T.M.B.), Department of Pediatrics and Physiology, University of Arkansas for Medical Sciences, Little Rock, Arkansas 72202; and Physics Department (G.I.L.), University of North Dakota, Grand Forks, North Dakota 58202

In a controlled feeding study, the effects of substituting 25 g soy protein for meat on calcium retention and bone biomarkers were determined. Postmenopausal women (n = 13) ate two diets that were similar, except that, in one diet, 25 g high-isoflavone soy protein (SOY) was substituted for an equivalent amount of meat protein (control diet), for 7 wk each in a randomized crossover design. After 3 wk of equilibration, calcium retention was measured by labeling the 2-d menu with ⁴⁵Ca, followed by whole-body counting for 28 d. Urinary calcium and renal acid excretion were measured at wk 3, 5, and 7. Biomarkers of bone and cardiovascular health were mea-

sured at the beginning and end of each diet. Calcium was similarly retained during the control and SOY diets (d 28, percent dose, mean \pm pooled SD: 14.1 and 14.0 \pm 1.6, respectively). Despite a 15–20% lower renal acid excretion during the SOY diet, urinary calcium loss was unaffected by diet. Diet also did not affect any of the indicators of bone or cardiovascular health. Substitution of 25 g high isoflavone soy protein for meat, in the presence of typical calcium intakes, did not improve or impair calcium retention or indicators of bone and cardiovascular health in postmenopausal women. (*J Clin Endocrinol Metab* 90: 181–189, 2005)

IN ADDITION TO providing structural framework and locomotion, the skeletal system serves as a buffering reservoir and aids the kidneys and lungs in the tight regulation of the systemic hydrogen ion concentration. Dietary practices that lead to chronic production of acid ash, such as diets high in meat, are hypothesized to tap into this alkali reservoir and cause a gradual dissolution of bone mineral (1, 2) and, as such, are considered a risk for hypercalciuria and osteoporosis (3–5). Although the sulfur amino acids in animal proteins, such as meat, are thought to cause hypercalciuria, the high phosphorus content of these proteins has been found to negate this effect (6). Many staple plant proteins, such as wheat and rice, have sulfur amino acid contents that are similar to or higher than those in meats (7), but the coexisting alkalis are thought to reduce the dietary acid load (8). Furthermore, the increased ammoniogenesis observed with higher protein intake may partly neutralize the acid production (9). Therefore, the net effect of a protein source on calcium balance is determined by many coexisting factors in the protein and in the whole diet and is therefore difficult to predict. Observational studies have indicated positive (10–12), negative (5, 13–15), or no association (16) between animal protein intake and bone health.

First Published Online October 13, 2004

Abbreviations: CONTROL, Control diet; GFR, glomerular filtration rate; HDL, high-density lipoprotein; IGF-IBP3, IGF-I binding protein-3; iPTH, intact PTH; LDL, low-density lipoprotein; NS, not significant; SOY, 25-g high-isoflavone soy protein (diet).

JCEM is published monthly by The Endocrine Society (<http://www.endo-society.org>), the foremost professional society serving the endocrine community.

Very few controlled feeding studies have tested the relative calciuric effect of animal *vs.* plant protein food sources, in the context of a mixed diet (17–19). In a short-term feeding study (12 d), under conditions of equalized phosphorus and low calcium (400 mg), higher net renal acid excretion and urinary calcium loss were observed from diets containing meat *vs.* soy protein (19). However, in a recent controlled feeding study of high- *vs.* low-meat diets of much longer duration (8 wk), in which phosphorus content of the diets was not manipulated and calcium content mimicked typical intakes (~600 mg/d), no increase in urinary calcium excretion or change in whole-body calcium retention was observed in postmenopausal women (20). Similar calcium retention occurred despite the higher sulfur amino acid intake and the higher urinary sulfate and renal acid excretion during the high-meat diet. Although these findings (20), combined with those from earlier controlled feeding studies (17, 18), indicate that a moderate increase in meat intake does not impair calcium retention, three short-term (3- to 6-month) supplementation studies suggest that incorporation of soy protein isolate in the diet may improve calcium homeostasis (21–23). Although these studies point to a component in soy protein isolate, such as isoflavones, as an enhancer of calcium absorption and/or utilization, the phytate and oxalate content of soy may reduce the intestinal absorption of calcium (24) and other minerals (25, 26); thus, the net effect of regular consumption of soy protein on calcium homeostasis is not known.

The primary objective of this carefully controlled feeding study was to determine the effects of daily substitution of

high-isoflavone, intact soy protein for meat protein, in a typical mixed diet, on calcium retention and bone metabolism in healthy postmenopausal women. Because the amount of soy protein (25 g) was selected to conform with the Food and Drug Administration (FDA)-approved health claim (27), the secondary objective was to test the effects of this dietary practice on indicators of cardiovascular health.

Subjects and Methods

Subjects

Postmenopausal women were recruited through public advertising (television, newspapers) and by direct mailings. The women were selected after an interview and blood analysis and qualified to enter the study if they were age 50–75 yr; at least 3 yr since last menses; had FSH more than 40 IU/liter; were nonsmokers; had no apparent underlying disease; had normal bone mineral density (femoral neck T score ≥ -2.5) as determined by dual-energy x-ray absorptiometry (Hologic Delphi QDR, Bedford, MA); had normal thyroid, liver, and kidney functions; were willing to discontinue any nutritional supplements as soon as their applications were received; and did not regularly use any medications (except for hormone replacement therapy).

The study was approved by the University of North Dakota Radioactive Drug Research Committee and Institutional Review Board, and

by the United States Department of Agriculture Radiological Safety Office. The study was explained verbally and in writing by the investigators, and written informed consent was given by each woman.

Of 15 women enrolled, two dropped out on the second day of the study (one because of transportation problems and the other without providing a reason). The remaining 13 (all white, six using hormone replacement therapy) were age 59.9 ± 5.0 yr (mean \pm SD; range, 52–69), with a body mass index of 26.0 ± 5.0 kg/m²; range, 19.7–38.5; median, 24.8; and femoral neck bone mineral density, 0.742 ± 0.093 (T-score range, –2.33 to –0.56). As estimated from 3-d food records, their calcium and protein intakes before the study were 857 ± 336 mg/d and 76 ± 23 g/d, respectively.

Diets

Registered dietitians planned two experimental diets using ordinary foods in a 2-d menu cycle (Table 1). The subjects consumed both diets for 7 wk, in a randomized crossover design, with a 2-wk break during which they consumed their self-selected diets but continued to take the multivitamin provided (Table 1). The weighed control diet (CONTROL) and 25-g high-isoflavone soy protein (SOY) diets provided (mean \pm SD) similar amounts of protein (15% of energy; 1.32 ± 0.19 and 1.33 ± 0.17 g/kg body weight) but contained 170 and 55 g/d meat, respectively (Table 1). The amount of calcium in the diet was chosen to be similar to typical intakes by postmenopausal women in the United States (28). Although the diets were planned with similar amounts of calcium, the

TABLE 1. Ingredient difference and nutrient composition of controlled high- and low-meat diets consumed by healthy postmenopausal women for 7 wk each in a crossover design^{a,b}

	CONTROL	SOY
Diet ingredient difference (g/d)		
d 1:		
Soy protein isolate (added to banana bread) ^c		+10
Mandarin oranges		+10
Beef ^d	+55	
Soy protein isolate (added to beef casserole) ^c		+10
Chicken ^d	+55	
White dinner roll		+5
Soy protein isolate (added to brownie) ^c		+10
d 2:		
Soy protein isolate (added to peanut butter) ^c		+5
Ham ^d	+60	
Soy protein isolate (added to gingerbread) ^c		+15
Chicken ^d	+60	
Soy protein isolate (added to cornbread) ^c		+10
Tub margarine	+1	
Nutrient composition		
Protein (% of energy)	15	15
Sulfur amino acids (g)	3.0	2.7
Fat (% of energy)	30	30
Saturated fat	7	7
Monounsaturated fat	13	13
Polyunsaturated fat	7	7
Cholesterol (mg)	220	157
Carbohydrate (% of energy)	55	55
Dietary fiber (g)	19	19
Calcium (mg)	690 (670 \pm 7)	746 (754 \pm 17)
Potassium (mg)	2617 (2584 \pm 290)	2303 (2587 \pm 339)
Sodium (mg)	3657 (3450 \pm 152)	3504 (3323 \pm 268)
Phosphorus (mg)	1505 (1596 \pm 87)	1533 (1638 \pm 104)
Magnesium (mg)	306	321
Phytate (mg)	1673	2246

^a All values are based on energy intake of 9.2 MJ (2200 kcal). Values for protein, carbohydrate, fat, cholesterol, and minerals were calculated from U.S. Department of Agriculture food composition data. Data in parentheses represent analyzed values for minerals (mean \pm SD).

^b A chewable multivitamin tablet containing 100% daily value (DV) for vitamins A, C, D, E, B₆, B₁₂, thiamin, riboflavin, niacin, folic acid, and pantothenic acid was taken daily by the participants.

^c The soy protein isolate contained 3.72 mg isoflavones/g protein (2.28 mg aglycone/mg protein; 1.22, 0.87, 0.19 mg/g protein of genistein, daidzein, and glycitein, respectively).

^d The protein content of beef, chicken, and ham was calculated from U.S. Department of Agriculture, Nutrient Database for Standard Reference, Release 14 (Nutrient Data Laboratory home page, <http://www.nal.usda.gov/fnic/foodcomp>) and was 22.03, 19.35, and 23.09 (g/100 g), respectively.

analyzed values for calcium were slightly higher in the SOY diet than the CONTROL (754 ± 17 *vs.* 670 ± 7 mg calcium/2200 kcal, respectively), reflecting the calcium content of the soy protein isolate. The isoflavone content of the isolate was 2.28 mg aglycone/G protein (product: IB1.2UN30CA, lot no. 038A-02; Solae Company, St. Louis, MO) (Table 1). Phytate content of the diets was calculated from food composition tables (29) (Table 1). To maintain body weights, energy intakes were adjusted by proportionally changing the amounts of all foods. The energy intakes were similar during the two dietary periods (2216 ± 271 and 2189 ± 300 kcal for CONTROL and SOY, respectively). Amounts of coffee, tea, and artificially sweetened, noncola, carbonated beverages (limited to two total servings daily), salt, and pepper were individualized and kept constant. City water and chewing gum were consumed as desired. Participants were given a list of approved over-the-counter medications and mouth products. The food items were prepared in oven-/microwave-oven-safe containers. Participants consumed the food quantitatively, using spatulas and rinse bottles, eating one meal at the research center on weekdays and the remaining foods elsewhere.

Calcium retention measurements with ^{47}Ca

Dietary calcium retention was measured with a ^{47}Ca radiotracer and whole-body scintillation counting (30) with adjustments to account for ^{47}Sc activity. The ^{47}Ca isotope was obtained by neutron activation (University of Missouri, Columbia, MO) of stable ^{46}Ca (as calcium carbonate, 30.89% enriched; Oak Ridge National Research Laboratory, Oakridge, TN). The custom-made scintillation counter (31) detects γ -emissions with 32 crystal NaI(Tl) detectors ($10 \times 10 \times 41$ cm each), arranged in two planes above and below a bed on which the subjects lie.

After 3 wk of dietary equilibration, all the meals in the 2-d menu were labeled with a total of 148 kBq ($4 \mu\text{Ci}$) ^{47}Ca ($<4 \mu\text{g}$ elemental calcium). Because of the concern that ingested calcium from some dietary sources may not form a common absorptive pool (32), both diets were designed with milk as the primary source of calcium. For each meal, the tracer was mixed with milk and allowed to equilibrate overnight. The specific activity (ratio of ^{47}Ca to elemental calcium) was constant for all meals for each individual during both dietary periods. The energy provided by the radiolabeled meals was constant during each 2-d administration. All labeled meals were consumed at the research center.

The initial total body activity was determined from the whole-body count, 1–3 h after the first labeled meal (before any isotope was excreted), divided by the fraction of the total activity that was in the first meal. Whole-body calcium retention was monitored for 28 d. Activity was corrected to the midpoint of the 2 d of labeled meals and adjusted for background and minor fluctuations in the measurement of a ^{47}Ca standard distributed in water (33). The precision of the whole-body counting measurements was 1.4%.

Analyses

The subjects provided total 48-h urine collections during wk 0, 3, 5, and 7 of each dietary period. Fasting blood samples were drawn at the beginning (wk 1) and end (wk 7) of each dietary period. For homocysteine and serum lipids, fasting blood samples were collected twice at the beginning and end of the study, separated by a few days, and the values were averaged. Calcium in the urine and acid-digested diet aliquots (34) was determined by inductively coupled argon plasma emission spectrophotometry. Mean (\pm SD) measurements were $98 \pm 4\%$ of certified values for calcium in a standard reference material (Typical Diet, 1548b, United States National Institute of Standards and Technology).

Urinary ammonium was determined colorimetrically (35) (Raichem, Hemagen Diagnostics, San Diego, CA). Titratable acidity was determined in undiluted urine by titrating to pH 7.40 with 0.1 N NaOH. Free organic acids were measured by the Van Slyke and Palmer (36) method as modified by Lemann *et al.* (37). Urinary sulfates were determined turbidometrically (38). ELISAs were used to determine bone-specific alkaline phosphatase (Metra Biosystems, Mountain View, CA) and estradiol (Abbott Laboratories, Abbott Park, IL). The inter- and intraassay variabilities were 5.2 and 5.0%, respectively, for bone-specific alkaline phosphatase and 6.2 and 6.7%, respectively, for estradiol. Serum tartrate-resistant acid phosphatase activity was determined using α -naphthylphosphate and diazotized-2-amino-5-chlorotoluene as substrates (39). Creatinine clearance was calculated from serum and urinary creatinine,

which were measured using alkaline picric acid (40). RIAs were used to determine serum TSH (TSH, T_3 , T_4) (Abbott Laboratories). The intra- and interassay variabilities were 4.2 and 5.4%, respectively, for TSH and 3.64 and 4.23%, respectively, for T_4 . RIAs were also used for intact PTH (iPTH), osteocalcin, and 25-hydroxyvitamin D (Diasorin, Stillwater, MN). The intra- and interassay variabilities were 3.6 and 3.4%, respectively, for iPTH; 4.3 and 11.9%, respectively, for osteocalcin; and 8.2 and 8.6%, respectively, for 25-hydroxyvitamin D. Serum IGF-I and IGF-I binding protein-3 (IGF-IBP3) (Diagnostic Systems Laboratory, Webster, TX) and urinary N-telopeptides (Ostex, Seattle, WA) were determined by ELISAs. The intra- and interassay variabilities were 7.1 and 5.4%, respectively, for IGF-I; 9.6 and 11.4% for IGF-IBP3; and 8.0 and 5.1% for N-telopeptides. Plasma ionized calcium was measured with an electrode (41) (8^+ Electrolyte Analyzer, Nova, Waltham, MA). The inter- and intraassay variabilities for this assay were 2.0 and 3.0%, respectively. Serum cholesterol fractions and triglycerides were determined using an automated procedure (Cobas Mira, Roche Diagnostic Systems, Inc., Sommerville, NJ). The inter- and intraassay variabilities were 1.0 and 1.2%, respectively, for total cholesterol; 2.1 and 2.6%, respectively, for high-density lipoprotein (HDL) fraction; and 1.0 and 3.0%, respectively, for triglycerides. Serum homocysteine was analyzed by fluorescence polarization immunoassay using an automated procedure (Abbott Laboratories). The inter- and intraassay variabilities were 2.2 and 5.2%, respectively.

Urine samples from wk 3 were analyzed for the unconjugated isoflavones (aglycones) and for total isoflavone content as previously described (42, 43). Briefly, for excreted aglycones, samples were extracted twice with 5 ml diethyl ether, and the organic layers were evaporated to dryness at 55 C under nitrogen. For conjugated isoflavones, samples were enzymatically hydrolyzed with *Helix pomatia* (sulfatase/glucuronidase, 100/1000 U) at 37 C for 3 h. All samples were then extracted twice with 5 ml diethyl ether, and the organic layers were evaporated to dryness at 55 C under nitrogen. Dried extracts were reconstituted in 0.5 ml of a solvent containing a known amount of biochanin A and injected into the LC-MS system to determine the aglycone concentrations using conditions reported previously (1, 2). All samples were analyzed in triplicate, and the results were expressed as nanograms/milligrams creatinine after normalization with biochanin A. Genistein (5,7,4'-trihydroxyisoflavone) and daidzein (7,4'-dihydroxyisoflavone) were purchased from Indofine Chemical Company, Inc. (Belle Mead, NJ). Sulfatase type V (aryl-sulfate sulfohydrolase) from *Helix pomatia* with reported sulfatase activity of 15–40 U/mg and glucuronidase activity of 400–600 U/mg was purchased from Sigma Chemical Company (St. Louis, MO).

Statistics

Individual ^{47}Ca retention data were modeled with a two-component exponential equation, $y = \beta_1 e^{-k_1 t} + \beta_2 e^{-k_2 t}$, where y represents ^{47}Ca retention as a percentage of the administered dose, t represents the time in hours, and coefficients β_1 and β_2 represent the fractional biological turnover of the radiotracer, expressed as percent of dose, at rates k_1 and k_2 , respectively. The calcium retention data for d 2–5 were not included in the model because they primarily represent the delay in elimination of the unabsorbed isotope. The percentage of ^{47}Ca initially absorbed was separately estimated from the y -intercept of the linear portion (d 9–25) of a semilogarithmic retention plot of percent ^{47}Ca retained *vs.* time.

Diet and sequence effects were evaluated for the parameter estimates in the calcium retention models, calcium retention on d 7, 14, 21, and 28, and initial calcium absorption. Urinary and blood measurements (wk 3, 5, and 7) were analyzed by using repeated-measures ANOVA followed by Tukey's contrasts (44). Variances in the data were expressed as pooled SD from the ANOVA. Sequence effects were evaluated before testing for diet and time effects and were found to be significant ($P < 0.1$) only for serum creatinine, homocysteine, low-density lipoprotein (LDL), total HDL, and HDL-3. For these variables, the mixed model was modified to use the diet \times time \times sequence interaction as the error term for the primary effects of diet and time. When data were highly skewed, they were logarithmically transformed so that the distribution would more closely approximate a normal distribution. For these, geometric means are reported in addition to the mean and pooled SD of the transformed data. The urinary isoflavone data were highly variable and were therefore ranked before ANOVA, and median and range of values are re-

ported. Using two-tailed probabilities, $P \leq 0.05$ was considered significant.

Results

Whole-body retention and intestinal absorption of calcium

Substitution of 25 g soy protein for meat protein did not affect the efficiency of calcium retention at any of the weekly time points tested (Table 2 and Fig. 1). By d 28, 14.0 and 14.1 (percent dose, ± 1.6) of the calcium tracer was retained from the CONTROL and SOY diets, respectively (Table 2, Fig. 1). As indicated by the two-component exponential model, with both diets, more than 75% of the tracer was eliminated rapidly (biological half-life, 1.5 d), indicating the excretion of the unabsorbed isotope and early endogenous losses. The remaining tracer ($\sim 20\%$) was eliminated less rapidly, with a biological half-life of 45–49 d (Table 2). Despite the higher phytate content of the SOY diet (by 573 mg/d), the initial absorption of calcium, expressed as percent dose, was also similar between the two diets [mean \pm pooled SD, 26.1 *vs.* 27.0% ± 7.0 ; not significant (NS); Table 2]. Calcium retention was not different between the two diets, at any time points tested, in women using hormone replacement therapy ($n = 6$).

Urine composition

Urinary pH was higher with the SOY, compared with CONTROL, by an average of about 0.15 pH units across all time points tested (overall mean, 6.33 *vs.* 6.19, respectively; $P \leq 0.0001$; Table 3) and decreased with time on both diets ($P = 0.002$). Titratable acidity was also lower by wk 3 of the SOY diet, and this difference was sustained throughout the diet period (overall mean, 41.5 *vs.* 53.4 mEq/d for SOY and CONTROL, respectively; $P = 0.03$; Table 3). Diet did not affect ammonium excretion at any of the time points tested (overall mean, 37.6 *vs.* 43.9 mEq/d for SOY and CONTROL, respectively, NS; Table 3), at least partially reflecting the

TABLE 2. Whole-body retention of a calcium radiotracer (^{47}Ca), measured by whole-body scintillation counting, as affected by daily substitution of SOY for an equivalent amount of CONTROL for 7 wk in healthy postmenopausal women^{a,b}

	CONTROL	SOY	Pooled SD	P
β_1 (% dose)	77.4	78.5	3.0	NS
β_2 (% dose)	22.7	21.6	2.9	NS
$\ln(T_1)$	0.43	0.43	0.09	NS
[Days] ^c	[1.5]	[1.5]		
$\ln(T_2)$	3.8	3.9	0.13	NS
[Days] ^c	[44.7]	[49.4]		
Retention @ d 7 (%)	24.4	23.5	4.8	NS
Retention @ d 14 (%)	18.2	17.7	2.3	NS
Retention @ d 21 (%)	15.9	15.6	1.7	NS
Retention @ d 28 (%)	14.1	14.0	1.6	NS
Absorption (% of dose) ^d	26.1	27.0	7.0	NS

^a Mean \pm SD, $n = 12$.

^b A two-component exponential model fit the data (R^2 of 0.97 to 0.99 for models of individuals on each diet). β_1 and β_2 represent the percentage of the isotope having a biological half-life of T_1 and T_2 , respectively. One subject's data could not be modeled using the two-component exponential equation.

^c Geometric means.

^d Intestinal calcium absorption, as percentage of dose, was estimated by using the linear portion (d 9–25 after ^{47}Ca administration) of a semilogarithmic retention plot [$\ln(\% \text{ retention } vs. \text{ time})$] and extrapolating back to the time of tracer administration.




FIG. 1. A, Whole-body calcium retention, as percent of dose (mean \pm SEM, $n = 13$) over time. Calcium retention was similar between the CONTROL and SOY diets at all weekly time points tested (see Table 2). B, The two component exponential models that fit the data for 12 of the 13 subjects. These group models ($y = 77.4 e^{-0.0196t} + 22.7 e^{-0.0007t}$ for high-meat and $y = 78.5 e^{-0.0191t} + 21.6 e^{-0.0007t}$ for CONTROL, where t is expressed in hours) use the mean coefficients from the modeled retention curves for each individual. Because of the variability in the early elimination of the isotope, calcium retention data for the first 2–5 d were omitted before the modeling of the data. The retention curves, modeled separately for individual on each diet, had coefficients of determination (R^2) of 0.97–0.99.

similar nitrogen content of the two menus. However, renal acid excretion, defined as the sum of ammonium and titratable acidity, was consistently lower during the SOY dietary period (overall mean, 97.2 *vs.* 79.1 mEq/d; $P = 0.0001$ for CONTROL and SOY diets, respectively; Table 3). Urinary free organic acid (representing compounds such as citric, acetic, and lactic acids) was similar during the two dietary periods. Urinary sulfate excretion was initially lower during the SOY than the CONTROL diet at wk 3, but this difference abated by wk 7 because of a gradual decrease during the CONTROL and an increase during the SOY dietary period (diet \times time interaction, $P = 0.01$, Table 3). Diet did not affect creatinine clearance measured at the end of each dietary period (1.33 *vs.* 1.27 ml/sec for CONTROL and SOY diets, respectively, NS; Table 3).

Despite the lower urine pH, and approximately 15–20% lower renal acid excretion during the SOY diet, urinary calcium excretion was similar between the two diets at all time points tested (overall mean, 3.53 and 3.48 mmol/d, NS, for the CONTROL and SOY diets, respectively; Table 3). No correlation between renal acid and urinary calcium excretion was detected. Urinary calcium, phosphorus, and oxalate excretion slightly increased, and urinary pH decreased, over time during both dietary periods ($P < 0.002$; Table 3).

TABLE 3. Urine composition of total 48-h composites as affected by a daily substitution of SOY for an equivalent amount of CONTROL for 7 wk in healthy postmenopausal women

	CONTROL			SOY			Pooled SD	Diet <i>P</i> value	Time <i>P</i> value
	Wk 3	Wk 5	Wk 7	Wk 3	Wk 5	Wk 7			
pH	6.25	6.17	6.13	6.41	6.29	6.30	0.12	0.0001	0.002
Titrateable acidity (mEq/d)	28.2	25.9	25.1	19.9	20.4	20.2	8.1	0.03	NS
Ammonium (mmol/d)	43.5	41.9	46.1	35.0	35.5	42.2	5.8	NS	NS
Renal acid excretion (mEq/d) ^a	70.9	66.3	69.4	55.0	56.4	62.3	10.5	0.0001	NS
ln (free organic acid) (mEq/d)	1.82 [6.17]	1.94 [6.96]	1.57 [4.81]	1.22 [3.39]	1.63 [5.10]	1.75 [5.75]	0.56	NS	NS
Sulfate (mmol/d) ^b	17.0 ^x	15.3 ^{x,y}	16.8 ^x	14.3 ^x	15.4 ^{x,y}	16.4 ^x	1.7	0.01	0.03
ln (calcium) [mg/d (mmol/d)]	4.86 (3.25) [130.0]	4.81 (3.09) [123.6]	5.06 (3.97) [158.8]	4.86 (3.22) [128.7]	4.88 (3.32) [132.6]	5.05 (3.93) [157.2]	0.22	NS	0.001
Phosphorus [mg/d (mmol/d)]	964.9 (31.2)	866.8 (28.0)	1167.5 (37.7)	874.4 (28.2)	896.8 (29.0)	1097.5 (35.4)	173.2	NS	0.0001
Oxalate [mg/d (μmol/d)]	26.2 (291)	24.5 (272)	28.3 (314)	21.4 (238)	24.2 (269)	27.5 (305)	3.6	NS	0.0004
Creatinine [g/d (mmol/d)]	1.10 (9.7)	1.11 (9.8)	1.14 (10.1)	0.98 (8.6)	1.01 (8.9)	1.07 (9.5)	0.12	0.003	NS
ln (N-telopeptide), nM BCE/mM creatinine			3.08 [21.8]			3.20 [25.0]	0.24	NS	
Hydroxyproline [mg/d (μmol/d)]			0.108 (107.9)			0.105 (104.5)	17	NS	
Creatinine clearance (ml/sec)			1.38			1.37	0.16	NS	

Data are means with pooled SD from ANOVA (*n* = 13). *P* values are from ANOVA. Data in brackets are geometric means.

^a Defined as sum of titrateable acidity and ammonium excretion. BCE, Bone collagen equivalents.

^b Significantly affected by diet × time interaction. Means without a common superscript (*x*, *y*) are significantly different as determined by Tukey's contrasts (*P* < 0.05).

Biomarkers of soy intake

Urinary isoflavone concentrations were used as a confirmatory marker of soy protein intake. The mean (\pm SEM) total urinary isoflavone concentrations were 100.41 ± 11.39 μ mol/mg creatinine in the soy-consuming women and 7.5 ± 1.30 μ mol/mg creatinine when no added soy protein was consumed. Based on our previous experience, the total urinary isoflavone excretion was consistent with the amount of soy consumed (42, 43). Furthermore, none of the women in this study were considered equal producers, because the mean equal excretion rate during the soy intake periods was 13.69 ± 4.46 μ mol/d (range, 1.22–9.49).

Biomarkers of bone metabolism

Substitution of soy protein for meat did not affect bone metabolism as indicated by specific blood biomarkers of bone formation (serum bone-specific alkaline phosphatase, osteocalcin, and IGF-I) or of bone resorption (serum tartrate-resistant acid phosphatase) (Table 4). Several other blood and urinary indices of bone and mineral metabolism, iPTH, 25-hydroxyvitamin D (Table 4), cortisol, plasma zinc, magnesium, calcium, phosphorus, ionized calcium, ionized magnesium (data not shown), and urinary N-telopeptide and hydroxyproline excretion, were also unaffected by dietary treatments (Table 3). Diet did not change any of the measured indicators of thyroid function (Table 4). The subjects were replete with vitamin D as indicated by serum 25-hydroxyvitamin D at wk 1 of both dietary periods (71.8 and 68.2 nmol/liter for CONTROL and SOY, respectively). Although this

study was conducted during October–March in Grand Forks, ND (latitude, 47.5° N), the vitamin D status of the subjects was successfully maintained and even slightly increased with a daily vitamin D₃ supplement of 20 μ g/d (Table 4).

Biomarkers of cardiovascular health

Daily substitution of soy protein for meat protein did not affect serum homocysteine concentration or the lipid profile (LDL, HDL, total cholesterol, and triacylglycerol) (Table 5). Diet also did not change any of the measured hemostatic indicators, such as fibrinogen, protime, and partial thromboplastin time (data not shown).

Discussion

Effects of meat *vs.* soy protein on calcium homeostasis

The results of this carefully controlled feeding study indicate that a daily substitution of 25 g intact, high isoflavone soy protein for an equivalent amount of meat protein for several weeks, in a mixed diet with typical calcium content, does not improve or impair calcium homeostasis in healthy postmenopausal women. This conclusion is supported by the whole-body calcium retention data (Table 2 and Fig. 1) and by a host of biomarkers of bone metabolism (Tables 3 and 4).

The design of this study was optimized for comparison of calcium retention by assuring sufficient statistical power (12 subjects were needed to detect a difference of 2.5 percentage points in calcium retention in a crossover design; 90% power; α = 0.05; residual SD = 1.5). The measurements were made

TABLE 4. Blood biomarkers of bone and mineral status as affected by daily substitution of SOY for an equivalent amount of CONTROL for 7 wk in healthy postmenopausal women

	CONTROL	SOY	Pooled SD	Diet <i>P</i> value	Time <i>P</i> value
Bone-specific alkaline phosphatase [U/liter (μ kat/liter)]					
Wk 1	17.9 (0.30)	18.4 (0.31)	1.9	NS	0.01
Wk 7	19.7 (0.33)	19.6 (0.33)			
ln (Osteocalcin) [ng/ml (nmol/liter)]	1.16 (–0.58) [3.20]	1.15 (–0.59) [3.16]	0.22	NS	NS
Serum IGF-I [ng/ml (nmol/liter)]	112.5 (14.8)	125.1 (16.8)	23.6	NS	NS
Serum IGF-IBP3 [ng/ml (nmol/liter)]	4711.3 (162.5)	4670.8 (161.1)	10.2	NS	NS
Tartrate-resistant acid phosphatase [U/liter (μ kat/liter)]	2.66 (44.3)	2.75 (45.8)	0.36	NS	NS
25-OH vitamin D [ng/ml (nmol/liter)]					
Wk 1	28.7 (71.8)	27.8 (68.2)	4.4	NS	0.004
Wk 7	30.5 (76.3)	33.8 (83.4)			
ln (iPTH) [pg/ml (pmol/liter)]	3.48 (1.25) [32]	3.49 (1.26) [33]	0.13	NS	NS
Total T ₃ [ng/ml (nmol/liter)]	71.0 (1.09)	72.9 (1.12)	6.5	NS	NS
Total T ₄ [μ g/dl (nmol/liter)]	8.8 (113.0)	9.1 (116.9)	0.6	NS	NS
TSH [μ U/ml (mU/liter)]	2.51 (2.51)	2.62 (2.62)	0.60	NS	NS

Data are mean with pooled SD (*n* = 13). *P* values are from ANOVA. The data in *brackets* are geometric means. Blood was drawn twice at the beginning and the end of each dietary period. When no time or diet \times time interaction was detected, the data from the four blood draws were pooled. μ kat, Microkatal.

after allowing for both equilibration and a break between the diets (total of 5 wk) and using sensitive radiotracer and whole-body counting methodology, in a crossover design. The subjects continued to consume the controlled diets during the whole-body counting measurements (4 additional wk); thus, the final retention data reflect not only the initial bioavailability of calcium but also the net effect of the subsequent loss through multiple excretory pathways. For this reason, the diets were designed to be similar except for the protein components, and the intakes of other factors known to influence calcium excretion (*e.g.* salt, potassium, caffeine) (28) were kept constant for each individual. The phosphorus content of the diets was similar without manipulation of the menus.

The hypercalciuric effect of protein is thought to be related to increased glomerular filtration rate (GFR) and reduced renal reabsorption of calcium in response to the acid-ash produced from sulfur amino acid catabolism (45–49). In the current study, despite lower urinary acidity and lower sulfate and renal acid excretions during the SOY diet, no change in creatinine clearance, a surrogate measure of GFR, was observed (Table 3). Therefore, it is not surprising that urinary calcium and whole-body calcium retention were not affected by the diets. This lack of responsiveness of urinary calcium excretion to both sulfate and renal acid excretions is consistent with our earlier findings (20). In a study of similar design, although renal acid excretion was about 45% higher

during wk 3 of the high-meat period (20), urinary calcium excretion was not different between controlled high- and low-meat diets (20% *vs.* 12% of energy as protein). This difference abated to only 18% by wk 8 of the study, indicating adaptation in renal acid excretion over time (20). These collective findings indicate that changes in urinary acidity within the range that may result from common, practical dietary practices do not reach a threshold that triggers an increase in GFR and/or use of body calcium as a buffering agent. Remer *et al.* (50) have suggested that a net acid excretion of more than 120 mEq/d is required for the depletion of plasma bicarbonate and use of alkali from the skeleton. In this study, the acid excretion (uncorrected for bicarbonate excretion) during the CONTROL diet (97 mEq/d) was below this threshold.

It is possible that diet-induced metabolic acidosis induces bone resorption through a cell-mediated mechanism (51), by inhibition of osteoblasts and stimulation of osteoclasts (52), without the direct involvement of the renal system. However, lack of changes in both bone resorption and formation biomarkers in this study do not support this hypothesis.

The current study provides strong evidence that a daily incorporation of soy protein (with high isoflavone content) in place of meat protein, in a mixed diet with typical calcium intakes of approximately 700 mg/d, does not improve calcium retention. However, several supplementation studies have suggested a bone sparing effect for high-isoflavone soy

TABLE 5. Serum homocysteine and lipid concentrations as affected by daily substitution of SOY for an equivalent amount of CONTROL for 7 wk in healthy postmenopausal women

	CONTROL	SOY	Pooled SD	Diet <i>P</i> value
Homocysteine [μ g/L (μ mol/liter)]	1162.7 (8.6)	1149.2 (8.5)	94.6	NS
Cholesterol [mg/dl (mmol/liter)]	244.1 (6.31)	240.9 (6.23)	12.8	NS
LDL [mg/dL (mmol/liter)]	141.9 (3.67)	138.4 (3.58)	0.09	NS
Total HDL [mg/dl (mmol/liter)]	82.0 (2.12)	82.4 (2.13)	6.6	NS
HDL-3 [mg/dl (mmol/liter)]	68.4 (1.77)	68.8 (1.78)	3.5	NS
ln (Triacylglycerols) [mg/dl (mmol/liter)]	4.54 (0.06) [93.7]	4.52 (0.04) [91.8]	0.14	NS

Data are mean with pooled SD (*n* = 13) and represent average of two blood draws at the beginning (wk 0, 1) and end of each dietary period (wk 6 and 7).

Data in *brackets* are geometric means. None of the variables were significantly affected by diet, time, or diet \times time interaction.

protein compared with milk proteins in peri- and postmenopausal women (22, 23, 53). Studies in ovariectomized rats have also indicated protection from estrogen deficiency bone loss in animals fed intact soy protein *vs.* casein (54–56). Although the results of the human studies cited above (22, 23, 53) must be interpreted with full recognition of their short duration and small sample size, the favorable effects of soy protein supplementation on bone are intriguing and imply improved calcium retention compared with isolated milk proteins. This difference may be related to the higher calciuric effect of milk-based proteins *vs.* soy protein. In a recent study, urinary calcium excretion increased by 33% from baseline in those receiving 40 g/d isolated milk-based proteins compared with no change in those receiving soy protein (23).

Alternatively, the putative beneficial effect of soy protein supplementation in the previous studies may be related to the higher combined doses of both soy protein (40 g/d) and supplemental calcium (650–1400 mg/d) used (22, 23, 53). The results of two recent studies (57, 58) suggest that calcium intake may modulate the effect of protein on the skeleton and that, at high intakes, dietary calcium and protein may synergistically interact to favorably affect bone mass. The mechanism for this interaction is not known but is thought to be at least partially related to an increase in the concentration of serum IGF-I, an osteotropic growth factor. This stimulatory effect on serum IGF-I has been observed with protein supplements alone (59) and with milk supplementation providing both protein and calcium (60, 61). A recent study has indicated that supplemental soy protein may be more potent in increasing serum IGF-I than milk-based proteins in postmenopausal women (23), suggesting that the source of protein may modify the protein effect on serum IGF-I. This notion is not supported by the findings of the current study, given that both soy and meat protein consumption failed to affect serum IGF-I. However, the study design was not optimal to test this idea, because the subjects were generally well-nourished, and their protein intake was not supplemented, closely mimicking their usual diets.

Effects of meat vs. soy protein on indicators of cardiovascular health

In this study, daily substitution of 25 g soy protein for meat for 7 wk did not improve serum lipids (Table 5) in mildly hypercholesterolemic women. Although the dose of soy protein conformed to the FDA-approved health claim (27), the lack of an effect was somewhat expected, because the fat composition of the diets was similar, and also the beneficial effects of soy protein are recently shown to be less than initially reported. The landmark meta-analysis of 38 studies (62), the basis of the health claim, predicted a decrease in total and LDL cholesterol concentrations of 9% and 13%, respectively, with an average intake of 47 g/d soy protein compared with casein. However, other studies have shown smaller effects, of 2–7% or no effect, of soy protein or soy products (63, 64) and a slower response in postmenopausal women (21). The hypocholesterolemic effect of soy protein is thought to be related to increased T_4 levels (65). In this study, thyroid function was unaffected by soy protein intake (Table

4). The small increase in free T_3 during the SOY dietary period is consistent with a previous report (66) but may not be of clinical consequence.

Plasma total homocysteine concentration has been shown to be an independent indicator of atherosclerotic disease (67). The results of a recent study indicated that, compared with casein, 30–50 g/d soy, added to self-selected diets, improves plasma total homocysteine concentrations in hyperlipidemic individuals (68). In this controlled diet study, soy protein did not affect plasma total homocysteine in mildly hypercholesterolemic women with normal homocysteine levels.

In summary, preliminary evidence from previous supplementation studies indicates improvements in bone and cardiovascular health from consuming soy protein compared with isolated milk proteins. In this carefully controlled feeding study, the calcium retention data, with supportive clinical chemistry, provide strong evidence that incorporation of soy protein in place of meat protein, in amounts currently recommended and combined with typical calcium intakes, provides no improvements or deterioration in calcium homeostasis or cardiovascular health. The potential synergistic effects between dietary protein and calcium, and the mechanisms of how protein source may modulate this interaction to affect bone health, should be rigorously investigated. Defining the relationship between dietary protein (source and quantity) and calcium metabolism will have important implications for nutrition policy and evidence-based advice for prevention of the growing problem of osteoporosis.

Acknowledgments

We are thankful for the invaluable assistance of the staff at Grand Forks Human Nutrition Research Center: Carol Zito; Emily Nielsen, R.N.; Suzy Thorsen, R.D.; Brenda Hanson, R.D.; Debbie Krause, R.D.; Angela Scheett, R.D.; Jackie Nelson; and Sandra Gallagher. We thank Dr. Steve Morris of Missouri Research Reactor for processing of the radio-tracer. We are deeply indebted to the study participants.

Received February 27, 2004. Accepted September 23, 2004.

Address all correspondence and requests for reprints to: Dr. Fariba K. Roughead, United States Department of Agriculture, Agricultural Research Service, Grand Forks Human Nutrition Research Center, Grand Forks, North Dakota 58202-9034. E-mail: froughea@gfhnrc.ars.usda.gov.

This work was primarily supported by the United States Department of Agriculture, with additional support from the North Dakota Beef Commission.

Mention of a trademark or proprietary product does not constitute a guarantee or warranty of the product by the United States Department of Agriculture and does not imply its approval to the exclusion of other products that may also be suitable. The United States Department of Agriculture, Agricultural Research Service, Northern Plains Area, is an equal opportunity/affirmative action employer, and all agency services are available without discrimination.

References

1. Wachman A, Bernstein DS 1968 Diet and osteoporosis. *Lancet* 1:958–959
2. Goto K 1918 Mineral metabolism in experimental acidosis. *J Biol Chem* 36: 355–376
3. Hegsted DM 2001 Fractures, calcium, and the modern diet. *Am J Clin Nutr* 74:571–573
4. Bunker VW 1994 The role of nutrition in osteoporosis. *Br J Biomed Sci* 51: 228–240

5. Sellmeyer DE, Stone KL, Sebastian A, Cummings SR 2001 A high ratio of dietary animal to vegetable protein increases the rate of bone loss and the risk of fracture in postmenopausal women. *Am J Clin Nutr* 73:118–122
6. Spencer H, Kramer L, Osis D 1988 Do protein and phosphorus cause calcium loss? *J Nutr* 118:657–660
7. Massey LK 2003 Dietary animal and plant protein and human bone health: a whole foods approach. *J Nutr* 133:862S–865S
8. Remer T 2000 Influence of diet on acid-base balance. *Semin Dial* 13:221–226
9. Remer T 2001 Influence of nutrition on acid-base balance—metabolic aspects. *Eur J Nutr* 40:214–220
10. Hannan MT, Tucker KL, Dawson-Hughes B, Cupples LA, Felson DT, Kiel DP 2000 Effect of dietary protein on bone loss in elderly men and women: the Framingham Osteoporosis Study. *J Bone Miner Res* 15:2504–2512
11. Chiu JF, Yang CY, Wang PW, Yao WJ, Hsieh CC 1997 Long term vegetarian diet and bone mineral density in postmenopausal Taiwanese women. *Calcif Tissue Int* 60:245–249
12. Munger RG, Cerhan JR, Chiu BC 1999 Prospective study of dietary protein intake and risk of hip fracture in postmenopausal women. *Am J Clin Nutr* 69:147–152
13. Feskanich D, Willett WC, Stampfer MJ, Colditz GA 1996 Protein consumption and bone fractures in women. *Am J Epidemiol* 143:472–479
14. Metz JA, Anderson JJ, Gallagher Jr PN 1993 Intakes of calcium, phosphorus, and protein, and physical-activity level are related to radial bone mass in young adult women. *Am J Clin Nutr* 58:537–542
15. Abelow BJ, Holford TR, Insogna KL 1992 Cross-cultural association between dietary animal protein and hip fracture: a hypothesis. *Calcif Tissue Int* 50:14–18
16. Tesar R, Notelovitz M, Shim E, Kawell G, Brown J 1992 Axial and peripheral bone density and nutrient intakes of postmenopausal vegetarian and omnivorous women. *Am J Clin Nutr* 56:699–704
17. Spencer H, Kramer L, Osis D, Norris C 1978 Effect of a high protein (meat) intake on calcium metabolism in man. *Am J Clin Nutr* 31:2167–2180
18. Hunt JR, Gallagher SK, Johnson LK, Lykken GI 1995 High- versus low-meat diets: effects on zinc absorption, iron status, and calcium, copper, iron, magnesium, manganese, nitrogen, phosphorus, and zinc balance in postmenopausal women. *Am J Clin Nutr* 62:621–632
19. Breslau NA, Brinkley L, Hill KD, Pak CY 1988 Relationship of animal protein-rich diet to kidney stone formation and calcium metabolism. *J Clin Endocrinol Metab* 66:140–146
20. Roughead ZK, Johnson LK, Lykken GI, Hunt JR 2003 Controlled high meat diets do not affect calcium retention or indices of bone status in healthy postmenopausal women. *J Nutr* 133:1020–1026
21. Baum JA, Teng H, Erdman Jr JW, Weigel RM, Klein BP, Persky VW, Freels S, Surya P, Bakhit RM, Ramos E, Shay NF, Potter SM 1998 Long-term intake of soy protein improves blood lipid profiles and increases mononuclear cell low-density-lipoprotein receptor messenger RNA in hypercholesterolemic, postmenopausal women. *Am J Clin Nutr* 68:545–551
22. Alekel DL, Germain AS, Peterson CT, Hanson KB, Stewart JW, Toda T 2000 Isoflavone-rich soy protein isolate attenuates bone loss in the lumbar spine of perimenopausal women. *Am J Clin Nutr* 72:844–852
23. Arjmandi BH, Khalil DA, Smith BJ, Lucas EA, Juma S, Payton ME, Wild RA 2003 Soy protein has a greater effect on bone in postmenopausal women not on hormone replacement therapy, as evidenced by reducing bone resorption and urinary calcium excretion. *J Clin Endocrinol Metab* 88:1048–1054
24. Heaney RP, Weaver CM, Fitzsimmons ML 1991 Soybean phytate content: effect on calcium absorption. *Am J Clin Nutr* 53:745–747
25. Hurrell RF, Juillat MA, Reddy MB, Lynch SR, Dassenko SA, Cook JD 1992 Soy protein, phytate, and iron absorption in humans. *Am J Clin Nutr* 56:573–578
26. Lonnerdal B, Jayawickrama L, Lien EL 1999 Effect of reducing the phytate content and of partially hydrolyzing the protein in soy formula on zinc and copper absorption and status in infant rhesus monkeys and rat pups. *Am J Clin Nutr* 69:490–496
27. Food and Drug Administration 1999 Food labeling: health claims: soy protein and coronary heart disease. 21 CFR Part 101. Federal Register 64:57700–57733.
28. Institute of Medicine, Food and Nutrition Board 1997 Dietary reference intakes for calcium, phosphorus, magnesium, vitamin D, and fluoride. Washington DC: National Academy of Science
29. Harland BF, Oberleas D 1987 Phytate in foods. *World Rev Nutr Diet* 52:235–259
30. Lykken GI 1983 A whole body counting technique using ultralow doses of ⁵⁹Fe and ⁶⁵Zn in absorption and retention studies in humans. *Am J Clin Nutr* 37:652–662
31. Lykken GI, Ong HS, Alkhatib HA, Harris TR, Momcilovic B, Penland JG 2000 Perquisite spin-off from twenty-two years of measuring background in the whole body counter steel room. *Ann NY Acad Sci* 904:267–270
32. Weaver CM, Heaney RP 1991 Isotopic exchange of ingested calcium between labeled sources. Evidence that ingested calcium does not form a common absorptive pool. *Calcif Tissue Int* 49:244–247
33. Wielopolski L, Cohn SH 1984 Application of the library least-squares analysis to whole-body counter spectra derived from an array of detectors. *Med Phys* 11:528–533
34. Analytical Methods Committee 1960 Methods of destruction of organic matter. *Analyst* 85:643–656
35. Tietz WN 1986 Textbook of clinical chemistry. Philadelphia: WB Saunders Company; 1575–1588
36. Van Slyke DD, Palmer WW 1920 Studies of acidosis. The titration of organic acids in urine. *J Biol Chem* 41:567–585
37. Lemann Jr J, Litzow JR, Lennon EJ 1966 The effects of chronic acid loads in normal man: further evidence for the participation of bone mineral in the defense against chronic metabolic acidosis. *J Clin Invest* 45:1608–1614
38. Lundquist P, Martensson J, Sorbo B, Ohman S 1980 Turbidimetry of inorganic sulfate, ester sulfate, and total sulfur in urine. *Clin Chem* 26:1178–1181
39. Shaw LM, Brummund W, Dorio RJ 1977 An evaluation of a kinetic acid phosphatase method. *Am J Clin Pathol* 68:57–62
40. Burtis C, Ashwood ER, eds. 1999 Tietz textbook of clinical chemistry. 3rd ed. Philadelphia: WB Saunders Company
41. Fogh-Anderson N, Christianson TF, Komarny L, Siggaard-Anderson O 1978 Measurement of free calcium ion in capillary blood and serum. *Clin Chem* 24:1545–1552
42. Shelnutt SR, Cimino CO, Wiggins PA, Badger TM 2000 Urinary pharmacokinetics of the glucuronide and sulfate conjugates of genistein and daidzein. *Cancer Epidemiol Biomarkers Prev* 9:413–419
43. Shelnutt SR, Cimino CO, Wiggins PA, Ronis MJ, Badger TM 2002 Pharmacokinetics of the glucuronide and sulfate conjugates of genistein and daidzein in men and women after consumption of a soy beverage. *Am J Clin Nutr* 76:588–594
44. SAS Institute Inc 1999 SAS/STAT user's guide, version 8. Cary, NC: SAS Institute, Inc
45. Allen LH, Oddoye EA, Margen S 1979 Protein-induced hypercalciuria: a longer term study. *Am J Clin Nutr* 32:741–749
46. Kim Y, Linkswiler HM 1979 Effect of level of protein intake on calcium metabolism and on parathyroid and renal function in the adult human male. *J Nutr* 109:1399–1404
47. Schuette SA, Zemel MB, Linkswiler HM 1980 Studies on the mechanism of protein-induced hypercalciuria in older men and women. *J Nutr* 110:305–315
48. Hegsted M, Schuette SA, Zemel MB, Linkswiler HM 1981 Dietary calcium and calcium balance in young men as affected by level of protein and phosphorus intake. *J Nutr* 111:553–562
49. Hegsted M, Linkswiler HM 1981 Long-term effects of level of protein intake on calcium metabolism in young adult women. *J Nutr* 111:244–251
50. Remer T, Dimitriou T, Manz F 2003 Dietary potential renal acid load and renal net acid excretion in healthy, free-living children and adolescents. *Am J Clin Nutr* 77:1255–1260
51. Kraut JA, Mishler DR, Singer FR, Goodman WG 1986 The effects of metabolic acidosis on bone formation and bone resorption in the rat. *Kidney Int* 30:694–700
52. Frick KK, Bushinsky DA 2003 Metabolic acidosis stimulates RANKL RNA expression in bone through a cyclo-oxygenase-dependent mechanism. *J Bone Miner Res* 18:1317–1325
53. Potter SM, Baum JA, Teng H, Stillman RJ, Shay NF, Erdman Jr JW 1998 Soy protein and isoflavones: their effects on blood lipids and bone density in postmenopausal women. *Am J Clin Nutr* 68(6 Suppl):1375S–1379S
54. Arjmandi BH, Alekel L, Hollis BW, Amin D, Stacewicz-Spuntzak M, Guo P, Kukreja SC 1996 Dietary soybean protein prevents bone loss in an ovariectomized rat model of osteoporosis. *J Nutr* 126:161–167
55. Arjmandi BH, Getlinger MJ, Goyal NV, Getlinger MJ, Juma S, Alekel L, Hasler CM, Kukreja SC 1998 Role of soy protein with normal or reduced isoflavone content in reversing bone loss induced by ovarian hormone deficiency in rats. *Am J Clin Nutr* 68(6 Suppl):1358S–1363S
56. Arjmandi BH, Birnbaum R, Goyal NV, Getlinger MJ, Juma S, Alekel L, Hasler CM, Drum M, Holis BW, Kukreja SC 1998 Bone-sparing effect of soy protein in ovarian hormone-deficient rats is related to its isoflavone content. *Am J Clin Nutr* 68(6 Suppl):1364S–1368S
57. Meyer HE, Pedersen JL, Loken EB, Verdal A 1997 Dietary factors and the incidence of hip fracture in middle-aged Norwegians. A prospective study. *Am J Epidemiol* 145:117–123
58. Dawson-Hughes B, Harris SS 2002 Calcium intake influences the association of protein intake with rates of bone loss in elderly men and women. *Am J Clin Nutr* 75:773–779
59. Bonjour JP, Schurch MA, Chevalley T, Ammann P, Rizzoli R 1997 Protein intake, IGF-I and osteoporosis. *Osteoporos Int* 7(Suppl 3):S36–S42
60. Cadogan J, Eastell R, Jones N, Barker ME 1997 Milk intake and bone mineral acquisition in adolescent girls: randomised, controlled intervention trial. *BMJ* 315:1255–1260
61. Heaney RP, McCarron DA, Dawson-Hughes B, Oparil S, Berga SL, Stern JS, Barr SI, Rosen CJ 1999 Dietary changes favorably affect bone remodeling in older adults. *J Am Diet Assoc* 99:1228–1233

62. Anderson JW, Johnstone BM, Cook-Newell ME 1995 Meta-analysis of the effects of soy protein intake on serum lipids. *N Engl J Med* 333:276–282
63. Gardner CD, Newell KA, Cherin R, Haskell WL 2001 The effect of soy protein with or without isoflavones relative to milk protein on plasma lipids in hypercholesterolemic postmenopausal women. *Am J Clin Nutr* 73:728–735
64. Wong WW, Smith EO, Stuff JE, Hachey DL, Heird WC, Pownell HJ 1998 Cholesterol-lowering effect of soy protein in normocholesterolemic and hypercholesterolemic men. *Am J Clin Nutr* 68(6 Suppl):1385S–1389S
65. Forsythe 3rd WA 1995 Soy protein, thyroid regulation and cholesterol metabolism. *J Nutr* 125(3 Suppl):619S–623S
66. Persky VW, Turyk ME, Wang L, Freels S, Chatterton R, Barnes S, Erdamn J, Sepkovic DW, Bradlow HL, Potter S 2002 Effect of soy protein on endogenous hormones in postmenopausal women. *Am J Clin Nutr* 75: 145–153
67. Gerhard GT, Duell PB 1999 Homocysteine and atherosclerosis. *Curr Opin Lipidol* 10:417–428
68. Tonstad S, Smerud K, Hoie L 2002 A comparison of the effects of 2 doses of soy protein or casein on serum lipids, serum lipoproteins, and plasma total homocysteine in hypercholesterolemic subjects. *Am J Clin Nutr* 76: 78–84

Erratum

In the article “Preoperative Evaluation of Infants with Focal or Diffuse Congenital Hyperinsulinism by Intravenous Acute Insulin Response Tests and Selective Pancreatic Arterial Calcium Stimulation” by Charles A. Stanley, Paul S. Thornton, Arupa Ganguly, Courtney MacMullen, Patricia Underwood, Pooja Bhatia, Linda Steinkrauss, Laura Wanner, Robin Kaye, Eduardo Ruchelli, Mariko Suchi, and N. Scott Adzick (*The Journal of Clinical Endocrinology & Metabolism* 89:288–296, 2004), there is an error in Table 5. The K_{ATP} channel mutation $K_{ir}6.2$ region should be listed as A101N instead of A102N. *The authors regret the error.*

JCEM is published monthly by The Endocrine Society (<http://www.endo-society.org>), the foremost professional society serving the endocrine community.